arXiv:1606.07534v1 [math.CV] 24 Jun 2016

Essential norms of Volterra-type operators between *Zygmund* type spaces*

Shanli Ye†
(*School of Sciences, Zhejiang University of Science and Technology, Hangzhou 310023, China*)
Caishu Lin
(*Department of Mathematics, Fujian Normal University, Fuzhou 350007, China*)

Abstract

In this paper, we investigate the boundedness of some Volterra-type operators between *Zygmund* type spaces. Then, we give the essential norms of such operators in terms of g, φ, their derivatives and the n-th power φ^n of φ.

Key words Essential norms, boundedness, *Volterra* type operators , Zygmund type spaces.

2010 MR Subject Classification 47B33, 47B38, 30H99.

1 Introduction

Let $D = \{z : |z| < 1\}$ be the open unit disk in the complex plane $\mathbf{C}$ and $\partial D = \{z : |z| = 1\}$ be its boundary, and $H(D)$ denote the set of all analytic functions on D.

For every $0 < \alpha < \infty$, we denote by $\mathcal{B}^\alpha$ the Bloch type space of all functions $f \in H(D)$ satisfying

$$b^\alpha(f) = \sup_{z\in D}(1 - |z|^2)^\alpha |f'(z)| < \infty$$

endowed with the norm $\|f\|_{\mathcal{B}^\alpha} = |f(0)| + b^\alpha(f)$. The little Bloch type space $\mathcal{B}_0^\alpha$ consists of all $f \in \mathcal{B}^\alpha$ satisfying $\lim_{|z|\to 1^-}(1 - |z|^2)^\alpha |f'(z)| = 0$, $\mathcal{B}_0^\alpha$ is obviously the closed subspace of $\mathcal{Z}^\alpha$. When $\alpha = 1$, we get the classical Bloch space $\mathcal{B}^1 = \mathcal{B}$ and little Bloch space $\mathcal{B}_0^1 = \mathcal{B}_0$. It is well-known that for $0 < \alpha < 1$, $\mathcal{B}^\alpha$ is a subspace of H^∞, the Banach space of bounded analytic functions on D.

For $0 < \alpha < \infty$ we denote by $\mathcal{Z}^\alpha$ the *Zygmund type space* of those functions $f \in H(D)$ satisfying

$$\sup_{z\in D}(1 - |z|^2)^\alpha |f''(z)| < \infty,$$

and the little Zymund type space $\mathcal{Z}_0^\alpha$ consists of all $f \in \mathcal{Z}^\alpha$ satisfying $\lim_{|z|\to 1^-}(1-|z|^2)^\alpha |f''(z)| = 0$, $\mathcal{Z}_0^\alpha$ is obviously the closed subspace of $\mathcal{Z}^\alpha$. It can easily proved that $\mathcal{Z}^\alpha$ is a Banach space under the norm

$$\|f\|_{\mathcal{Z}^\alpha} = |f(0)| + |f'(0)| + \sup_{z\in D}(1 - |z|^2)^\alpha |f''(z)| \tag{1.1}$$

*The research was supported by the National Natural Science Foundation of China (Grant No. 11571217) and the Natural Science Foundation of Fujian Province, China(Grant No. 2015J01005).

†Corresponding author. E-mail address: ye_shanli@aliyun.com; slye@zust.edu.cn

and that $\mathcal{Z}_0^\alpha$ is a closed subspace of $\mathcal{Z}^\alpha$. When $\alpha = 1$, we get the classical Zygmund space $\mathcal{Z}^1 = \mathcal{Z}$ and the little Zygmund space $\mathcal{Z}_0^1 = \mathcal{Z}_0$. It is clear that $f \in \mathcal{Z}$ if and only if $f' \in \mathcal{B}^1$.

We consider the weighted Banach spaces of analytic functions

$$H_v^\infty = \{f \in H(D) : \|F\|_v = \sup_{z\in D} v(z)|f(z)| < \infty\}$$

endowed with norm $\|\cdot\|_v$, where the weight $v : D \to R_+$ is a continuous, strictly positive and bounded function. The weight v is called radial, if $v(z) = v(|z|)$ for all $z \in D$. For a weight v the associated weight $\widetilde{v}$ is definded by

$$\widetilde{v}(z) = (\sup\{|f(z)| : f \in H_v^\infty, \|f\|_v \le 1\}^{-1}, \qquad z \in D.$$

We notice the standard weights $v_\alpha(z) = (1 - |z|^2)^\alpha$, where $0 < \alpha < \infty$ and it is well-known that $\widetilde{v_\alpha} = v_\alpha$. We also consider the logarithmic weight

$$v_{\log} = (\log \frac{2}{1 - |z|^2})^{-1}, \qquad z \in D.$$

It is straightforward to show that $\widetilde{v_{\log}} = v_{\log}$.

Let u be a fixed analytic function on D and an analytic self-map $\varphi : D \to D$. Define a linear operator uC_φ on the space of analytic functions on D, called a weighted composition operator, by $uC_\varphi f = u \cdot (f \circ \varphi)$, where f is an analytic function on D. It is interesting to provide a function theoretic characterization when φ and u induces a bounded or compact composition operator on various function spaces. Some results on the boundedness and compactness of concrete operators between some spaces of analytic functions one of which is of Zygmund-type can be found, e.g., in ([1, 4, 5, 6, 11, 12, 13, 14, 18, 19, 21]).

Suppose that $g : D \to \mathbb{C}$ is a analytic map. Let U_g and V_g denote the Volterra-type operators with the analytic symbol g on D respectively:

$$U_g f(z) = \int_0^z f(\xi)g'(\xi)d\xi, \qquad z \in D.$$

and

$$V_g f(z) = \int_0^z f'(\xi)g(\xi)d\xi, \qquad z \in D.$$

If $g(z) = z$, then U_g is an integral operator. While $g(z) = \ln \frac{1}{1-z}$, then U_g is Cesàro operator. Pommerenke introduced the $Volterra$ type operator U_g and characterized the boundedness of U_g between H^2 spaces in [10] . More recently, boundedness and compactness of $Volterra$ type operators between several spaces of analytic functions have been studied by many authors, one may see in [17, 20].

In this paper, we consider the following integral-type operators, which were introduced by Li and Stevi̇c (see e. g. [5, 7]), they can be defined by:

$$(C_\varphi U_g f)(z) = \int_0^{\varphi(z)} f(\xi)g'(\xi)d\xi, \qquad (C_\varphi V_g f)(z) = \int_0^{\varphi(z)} f'(\xi)g(\xi)d\xi,$$

$$(U_g C_\varphi f)(z) = \int_0^z f(\varphi(\xi))g'(\xi)d\xi, \qquad (V_g C_\varphi f)(z) = \int_0^z f'(\varphi(\xi))g(\xi)d\xi.$$

We will characterize the boundedness of those integral-type operators between Zygmund type spaces, and also estimate their essential norms. The boundedness and compactness of these operators on the logarithmic Bloch space have been characterized in [20].

Recall that essential norm $\|T\|_{e,X\to Y}$ of a bounded linear operator $T : X \to Y$ is defined as the distance from T to $\mathbf{K}(X, Y)$, the space of compact operators from X to Y, namely,

$$\|T\|_{e,X\to Y} = \inf\{\|T + K\|_{X\to Y} : K : X \to Y is \ \ compact\}.$$

It provides a measure of non-compactness of T. Clearly, T is compact if and only if $\|T\|_{e,X\to Y} = 0$.

Throughout this paper, constants are denoted by C, they are positive and may differ from one occurrence to the other. The notation $a \asymp b$ means that there are positive constants C_1, C_2 such that $C_1 a \le b \le C_2 a$.

2 Boundedness

In order to prove the main results of this paper. we need some auxiliary results.

Lemma 2.1 ([1, 12]) *For $0 < \alpha < 2$ and let $\{f_n\}$ be a bounded sequence in $\mathcal{Z}^\alpha$ which converges to 0 uniformly on compact subsets of D . Then $\lim_{n\to\infty} \sup_{z\in D} |f_n(z)| = 0$.*

Lemma 2.2 ([1, 12]) *For every $f \in \mathcal{Z}^\alpha$ where $\alpha > 0$ we have:*
(i) $|f'(z)| \le \frac{2}{1-\alpha}\|f\|_{\mathcal{Z}^\alpha}$ and $|f(z)| \le \frac{2}{1-\alpha}\|f\|_{\mathcal{Z}^\alpha}$ for every $0 < \alpha < 1$;
(ii) $|f'(z)| \le 2\log\frac{2}{1-|z|}\|f\|_{\mathcal{Z}^\alpha}$ and $|f(z)| \le \|f\|_{\mathcal{Z}^\alpha}$ for $\alpha = 1$;
(iii) $|f'(z)| \le \frac{2}{1-\alpha}\frac{\|f\|_{\mathcal{Z}^\alpha}}{(1-|z|)^{\alpha-1}}$, for every $\alpha > 1$;
(iv) $|f(z)| \le \frac{2}{(\alpha-1)(2-\alpha)}\|f\|_{\mathcal{Z}^\alpha}$, for every $1 < \alpha < 2$;
(v) $|f(z)| \le 2\log\frac{2}{1-|z|}\|f\|_{\mathcal{Z}^\alpha}$, for every $\alpha = 2$;
(vi) $|f(z)| \le \frac{2}{(\alpha-1)(\alpha-2)}\frac{\|f\|_{\mathcal{Z}^\alpha}}{(1-|z|)^{\alpha-2}}$, for every $\alpha > 2$.

Lemma 2.3 ([1]) *Let $0 < \alpha < \infty$, v a radial, non-increasing weights tending to 0 at boundary of D and the weighted composition operator $uC_\varphi : \mathcal{Z}_\alpha \to H^\infty_v$ be bounded.*
(i) If $0 < \alpha < 2$, then uC_φ is a compact operator.
(ii)

$$\begin{aligned}\|uC_\varphi\|_{e,\mathcal{Z}_2\to H^\infty_v} &\asymp \limsup_{n\to\infty}(\log n)\|u\varphi^n\|_v \\ &\asymp \limsup_{|\varphi(z)|\to 1} v(z)|u(z)|\log\frac{2}{1-|\varphi(z)|^2}.\end{aligned}$$

(iii) If $\alpha > 2$, then

$$\begin{aligned}\|uC_\varphi\|_{e,\mathcal{Z}_\alpha\to H^\infty_v} &\asymp \limsup_{n\to\infty}(n+1)^{\alpha-2}\|u\varphi^n\|_v \\ &\asymp \limsup_{|\varphi(z)|\to 1}\frac{v(z)|u(z)|}{(1-|\varphi(z)|^2)^{\alpha-2}}.\end{aligned}$$

The following lemma is due to [2] and [8].

Lemma 2.4 *Let v and w be radial, non-increasing weights tending to zero at the boundary of D. Then*

(i) The weighted composition operator uC_φ maps H^∞_v into H^∞_w if and only if

$$sup_{n\geq 0}\frac{\|u\varphi^n\|_w}{\|z^n\|_v} \asymp \sup_{z\in D}\frac{w(z)}{\widetilde{v}(\varphi(z))}|u(z)| < \infty,$$

with norm comparable to the above supermum.

(ii)

$$\|uC_\varphi\|_{e,H^\infty_v\to H^\infty_w} = \limsup_{n\to\infty}\frac{\|u\varphi^n\|_w}{\|z^n\|_v} = \limsup_{|\varphi(z)|\to 1}\frac{w(z)}{\widetilde{v}(\varphi(z))}|u(z)|.$$

Lemma 2.5 ([3]) *For every $0<\alpha<\infty$, we have*

$$\lim_{n\to\infty}(n+1)^\alpha\|z^n\|_{v_\alpha} = (\frac{2\alpha}{e})^\alpha \tag{2.1}$$

and

$$\lim_{n\to\infty}(\log n)\|z^n\|_{v_{\log}} = 1. \tag{2.2}$$

Theorem 2.1 *Let φ be an analytic self-map of D and $g\in H(D)$.*

(i) If $0<\alpha<1$, then $V_gC_\varphi : \mathcal{Z}^\alpha\to\mathcal{Z}^\beta$ is a bounded operator if and only if $g'\in H^\infty_{v_\beta}$ and

$$\sup_{n\geq 0}(n+1)^\alpha\|g(\varphi')\varphi^n\|_{v_\beta} \asymp \sup_{z\in D}\frac{(1-|z|^2)^\beta}{(1-|\varphi(z)|^2)^\alpha}|g(z)\varphi'(z)| < \infty. \tag{2.3}$$

(ii) If $\alpha=1$, then $V_gC_\varphi : \mathcal{Z}^\alpha\to\mathcal{Z}^\beta$ is a bounded operator if and only if

$$\sup_{n\geq 0}(n+1)\|g(\varphi')\varphi^n\|_{v_\beta} \asymp \sup_{z\in D}\frac{(1-|z|^2)^\beta}{1-|\varphi(z)|^2}|g(z)\varphi'(z)| < \infty \tag{2.4}$$

and

$$\sup_{n\geq 0}(\log n)\|g'\varphi^n\|_{v_\beta} \asymp \sup_{z\in D}(1-|z|^2)^\beta\log(\frac{2}{1-|\varphi(z)|^2})|g'(z)| < \infty. \tag{2.5}$$

(iii) If $\alpha>1$, then $V_gC_\varphi : \mathcal{Z}^\alpha\to\mathcal{Z}^\beta$ is a bounded operator if and only if

$$\sup_{n\geq 0}(n+1)^\alpha\|g(\varphi')\varphi^n\|_{v_\beta} \asymp \sup_{z\in D}\frac{(1-|z|^2)^\beta}{(1-|\varphi(z)|^2)^\alpha}|g(z)\varphi'(z)| < \infty \tag{2.6}$$

and

$$\sup_{n\geq 0}(n+1)^{\alpha-1}\|g'\varphi^n\|_{v_\beta} \asymp \sup_{z\in D}\frac{(1-|z|^2)^\beta}{(1-|\varphi(z)|^2)^{\alpha-1}}|g'(z)| < \infty. \tag{2.7}$$

Proof Suppose that V_gC_φ is bounded from $\mathcal{Z}^\alpha$ to $\mathcal{Z}^\beta$. Using the test functions $f(z)=z$ and $f(z)=z^2$, we have

$$(1-|z|^2)^\beta|(V_gC_\varphi z)''| = (1-|z|^2)^\beta|g'(z)| < \infty,$$

and

$$(1-|z|^2)^\beta|(V_gC_\varphi z^2)''| = (1-|z|^2)^\beta|2\varphi'(z)g(z)+2\varphi(z)g'(z)| < \infty.$$

Since φ is a self-map, we get that $g'\in H^\infty_{v_\beta}$, $\varphi' g\in H^\infty_{v_\beta}$.

For every $0<\alpha<\infty$ and given nonzero $a\in D$. We take the test functions

$$f_a(z) = \frac{1}{\overline{a}}[\frac{(1-|a|^2)^2}{(1-\overline{a}z)^\alpha\alpha} - \frac{1-|a|^2}{(1-\overline{a}z)^{\alpha-1}}], \tag{2.8}$$

$$h_a(z) = \frac{1}{\overline{a}} \int_0^z \frac{1-|a|^2}{(1-\overline{a}w)^\alpha} dw, \tag{2.9}$$

$$g_a(z) = f_a(z) - h_a(z), \tag{2.10}$$

for every $z \in D$. One can show that f_a, h_a, and g_a are in $\mathcal{Z}_0^\alpha$, $\sup_{\frac{1}{2}<a<1} \|f_a\|_{\mathcal{Z}^\alpha} < \infty$ and $\sup_{\frac{1}{2}<a<1} \|h_a\|_{\mathcal{Z}^\alpha} < \infty$. Since $g_a'(a) = 0$, $g_a''(a) = \frac{\alpha}{(1-|a|^2)^\alpha}$, it follows that for all $z \in D$ with $|\varphi(z)| > \frac{1}{2}$, we have

$$\begin{aligned} +\infty > C\|g_a\|_{\mathcal{Z}^\alpha} &\geq \|V_g C_\varphi(g_{\varphi(z)})\|_{\mathcal{Z}^\beta} \\ &\geq (1-|z|^2)^\beta |\varphi'(z)g(z)||g''_{\varphi(z)}(\varphi(z))| - (1-|z|^2)^\beta |g'(z)||g'_{\varphi(z)}(\varphi(z))| \\ &= (1-|z|^2)^\beta |\varphi'(z)g(z)||\frac{\alpha}{(1-|\varphi(z)|^2)^\alpha}|. \end{aligned}$$

Then

$$\begin{aligned} \sup_{z\in D} |\varphi'(z)g(z)| \frac{(1-|z|^2)^\beta}{(1-|\varphi(z)|^2)^\alpha} &\leq \sup_{|\varphi(z)|\leq\frac{1}{2}} \frac{|\varphi'(z)g(z)|(1-|z|^2)^\beta}{(1-|\varphi(z)|^2)^\alpha} + \sup_{|\varphi(z)|>\frac{1}{2}} \frac{|\varphi'(z)g(z)|(1-|z|^2)^\beta}{(1-|\varphi(z)|^2)^\alpha} \\ &\leq (\frac{3}{4})^\alpha \|\varphi g'\|_{v_\beta} + C\|g_a\|_{\mathcal{Z}^\alpha} < \infty. \end{aligned}$$

Now we use (2) and Lemma 2.4 to conclude that

$$\sup_{n\geq 0} (n+1)^\alpha \|g(\varphi')\varphi^n\|_{v_\beta} \asymp \sup_{z\in D} \frac{(1-|z|^2)^\beta}{(1-|\varphi(z)|^2)^\alpha} |g(z)\varphi'(z)| < \infty,$$

which shows that (4) is necessary for all case.

Conversely, suppose that $g' \in H^\infty_{v_\beta}$ and (4) hold. Assume that $f \in \mathcal{Z}^\alpha$. From Lemma 2.2, it follows that

$$\begin{aligned} (1-|z|^2)^\beta |(V_g C_\varphi f)''(z)| &= (1-|z|^2)^\beta ||\varphi'(z)g(z)f''(\varphi(z)) + g'(z)f'(\varphi(z))| \\ &\leq (1-|z|^2)^\beta |\varphi'(z)g(z)f''(\varphi(z))| + (1-|z|^2)^\beta |g'(z)f'(\varphi(z))| \\ &\leq \frac{(1-|z|^2)^\beta}{(1-|\varphi(z)|^2)^\alpha} |\varphi'(z)g(z)| \|f\|_{\mathcal{Z}^\alpha} + C(1-|z|^2)^\beta |g'(z)| \|f\|_{\mathcal{Z}^\alpha} \\ &\leq C\|f\|_{\mathcal{Z}^\alpha}, \end{aligned}$$

$$|V_g C_\varphi(f)(0)| = 0, \quad \text{and} \qquad |(V_g C_\varphi f)'(0)| = |f'(\varphi(0))g(0)| \leq \|f\|_{\mathcal{Z}^\alpha} |g(\varphi(0))|,$$

which implies that $V_g C_\varphi$ is bounded. This completes the proof of (i).

Next we will prove (ii). The necessity in condition (5) has been proved above. Fixed $a \in D$ with $|a| > \frac{1}{2}$, we take the function

$$k_a(z) = \frac{p(\overline{a}z)}{\overline{a}} (\log \frac{1}{1-|a|})^{-1}, \tag{2.11}$$

for $z \in D$, where

$$p(z) = (z-1)((1+\log \frac{1}{1-z})^2 + 1).$$

Then we have $\sup_{\frac{1}{2}<|a|<1}\|k_a\|_{\mathcal{Z}^\alpha}\leq C$ by ([6]) . Let $a=\varphi(z)$, it follows that

$$\begin{aligned}\|V_gC_\varphi(k_{\varphi(z)})\|_{\mathcal{Z}^\beta} &\geq (1-|z|^2)^\beta|g'(z)||k'_{\varphi(z)}(\varphi(z))|-(1-|z|^2)^\beta|\varphi'(z)g(z)||k''_{\varphi(z)}(\varphi(z))|\\ &= (1-|z|^2)^\beta|g'(z)|\log\frac{1}{1-|\varphi(z)|^2}-(1-|z|^2)^\beta|\varphi'(z)g(z)||\frac{\alpha}{(1-|\varphi(z)|^2)^\alpha}|.\end{aligned}$$

Since (5) holds and V_gC_φ is bounded, we obtain that

$$\begin{aligned}\sup_{|\varphi(z)|>\frac{1}{2}}(1-|z|^2)^\beta\log\frac{1}{1-|\varphi(z)|^2}|g'(z)| &\leq \sup_{|\varphi(z)|>\frac{1}{2}}(1-|z|^2)^\beta|\varphi'(z)g(z)||\frac{\alpha}{(1-|\varphi(z)|^2)^\alpha}|\\ &+ \sup_{|\varphi(z)|>\frac{1}{2}}\|V_gC_\varphi(k_{\varphi(z)})\|_{\mathcal{Z}^\beta}\\ &< \infty.\end{aligned}$$

Noting $g'\in H^\infty_{v_\beta}$, and together with (3) and Lemma 2.4, we conclude that (6) holds.

The converse implication can be shown as in the proof of (i).

Finally we will prove (iii). We have proved that (7) holds above. To prove (8), we take function $f_{\varphi(z)}$ defined in (9) for every $z\in D$ with $|\varphi(z)|>\frac{1}{2}$, and obtain that

$$\begin{aligned}\|V_gC_\varphi(f_{\varphi(z)})\|_{\mathcal{Z}^\beta} &\geq (1-|z|^2)^\beta|g'(z)||f'_{\varphi(z)}(\varphi(z))|-(1-|z|^2)^\beta|\varphi'(z)g(z)||f''_{\varphi(z)}(\varphi(z))|\\ &= (1-|z|^2)^\beta|g'(z)|\frac{1}{|\varphi(z)|(1-|\varphi(z)|^2)^{\alpha-1}}\\ &- (1-|z|^2)^\beta|\varphi'(z)g(z)||\frac{2\alpha}{(1-|\varphi(z)|^2)^\alpha}|.\end{aligned}$$

Since V_gC_φ is bounded and (7) holds, we obtain that

$$\begin{aligned}\sup_{|\varphi(z)|>\frac{1}{2}}\frac{(1-|z|^2)^\beta}{(1-|\varphi(z)|^2)^{\alpha-1}}|g'(z)| &\leq \sup_{|\varphi(z)|>\frac{1}{2}}(1-|z|^2)^\beta|\varphi'(z)g(z)||\frac{2\alpha}{(1-|\varphi(z)|^2)^\alpha}|\\ &+ \sup_{|\varphi(z)|>\frac{1}{2}}\|V_gC_\varphi(f_{\varphi(z)})\|_{\mathcal{Z}^\beta}<\infty,\end{aligned}$$

therefore, we deduce that (8) holds by (2) and Lemma 2.4.

The converse implication can be shown as in the proof of (i).

Theorem 2.2 *Let φ be an analytic self-map of D and $g\in H(D)$.*

(i) If $0<\alpha<1$, then $C_\varphi V_g:\mathcal{Z}^\alpha\to\mathcal{Z}^\beta$ is a bounded operator if and only if $((g\circ\varphi)(\varphi'')+(g'\circ\varphi)(\varphi')^2)\in H^\infty_{v_\beta}$ and

$$\sup_{n\geq0}(n+1)^\alpha\|(g\circ\varphi)(\varphi')^2\varphi^n\|_{v_\beta}\asymp\sup_{z\in D}\frac{(1-|z|^2)^\beta}{(1-|\varphi(z)|^2)^\alpha}|g(\varphi(z))(\varphi'(z))^2|<\infty. \tag{2.12}$$

(ii) If $\alpha=1$, then $C_\varphi V_g:\mathcal{Z}^\alpha\to\mathcal{Z}^\beta$ is a bounded operator if and only if

$$\sup_{n\geq0}(n+1)\|(g\circ\varphi)(\varphi')^2\varphi^n\|_{v_\beta}\asymp\sup_{z\in D}\frac{(1-|z|^2)^\beta}{1-|\varphi(z)|^2}|g(\varphi(z))(\varphi'(z))^2|<\infty \tag{2.13}$$

and

$$\sup_{n\geq 0}(\log n)\|((g'\circ\varphi)(\varphi')^2+(g\circ\varphi)(\varphi''))\varphi^n\|_{v_\beta}\asymp\sup_{z\in D}(1-|z|^2)^\beta\log\frac{2}{1-|\varphi(z)|^2}\times|g'(\varphi(z))(\varphi'(z))^2+(g(\varphi(z))\varphi''(z)|<\infty. \tag{2.14}$$

(iii) If $\alpha>1$, then $C_\varphi V_g:\mathcal{Z}^\alpha\to\mathcal{Z}^\beta$ is a bounded operator if and only if (13) holds and

$$\sup_{n\geq 0}(n+1)^{\alpha-1}\|((g'\circ\varphi)(\varphi')^2+(g\circ\varphi)(\varphi''))\varphi^n\|_{v_\beta}\asymp\sup_{z\in D}\frac{(1-|z|^2)^\beta}{(1-|\varphi(z)|^2)^{\alpha-1}}\times|g'(\varphi(z))(\varphi'(z))^2+(g(\varphi(z))\varphi''(z)|<\infty. \tag{2.15}$$

The proof is similar to that of Theorem 2.1, the details are omitted.

Theorem 2.3 *Let φ be an analytic self-map of D and $g\in H(D)$.*

(i) If $0<\alpha<1$, then $C_\varphi U_g:\mathcal{Z}^\alpha\to\mathcal{Z}^\beta$ is a bounded operator if and only if $((g'\circ\varphi)(\varphi'')+(g''\circ\varphi)(\varphi')^2)\in H^\infty_{v_\beta}$ and $(g'\circ\varphi)(\varphi')^2\in H^\infty_{v_\beta}$.

(ii) If $\alpha=1$, then $C_\varphi U_g:\mathcal{Z}^\alpha\to\mathcal{Z}^\beta$ is a bounded operator if and only if $((g'\circ\varphi)(\varphi'')+(g''\circ\varphi)(\varphi')^2)\in H^\infty_{v_\beta}$ and

$$\sup_{n\geq 0}(\log n)\|(g'\circ\varphi)(\varphi')^2\varphi^n\|_{v_\beta}\asymp\sup_{z\in D}(1-|z|^2)^\beta\log(\frac{2}{1-|\varphi(z)|^2})|g'(\varphi(z))(\varphi'(z))^2|<\infty, \tag{2.16}$$

(iii) If $1<\alpha<2$, then $C_\varphi U_g:\mathcal{Z}^\alpha\to\mathcal{Z}^\beta$ is a bounded operator if and only if $((g'\circ\varphi)(\varphi'')+(g''\circ\varphi)(\varphi')^2)\in H^\infty_{v_\beta}$ and

$$\sup_{n\geq 0}(n+1)^{\alpha-1}\|((g'\circ\varphi)(\varphi')^2)\varphi^n\|_{v_\beta}\asymp\sup_{z\in D}\frac{(1-|z|^2)^\beta}{(1-|\varphi(z)|^2)^{\alpha-1}}\times|g'(\varphi(z))(\varphi'(z))^2|<\infty. \tag{2.17}$$

(iv) If $\alpha=2$, then $C_\varphi U_g:\mathcal{Z}^\alpha\to\mathcal{Z}^\beta$ is a bounded operator if and only if

$$\sup_{n\geq 0}(n+1)\|((g'\circ\varphi)(\varphi')^2)\varphi^n\|_{v_\beta}\asymp\sup_{z\in D}\frac{(1-|z|^2)^\beta}{(1-|\varphi(z)|^2)}\times|g'(\varphi(z))(\varphi'(z))^2|<\infty \tag{2.18}$$

and

$$\sup_{n\geq 0}(\log n)\|((g''\circ\varphi)(\varphi')^2+(g'\circ\varphi)\varphi'')\varphi^n\|_{v_\beta}\asymp\sup_{z\in D}(1-|z|^2)^\beta\log(\frac{2}{(1-|\varphi(z)|^2)})\times|g''(\varphi(z))(\varphi'(z))^2+g'(\varphi(z))\varphi''(z)|<\infty. \tag{2.19}$$

(v) If $\alpha>2$, then $C_\varphi U_g:\mathcal{Z}^\alpha\to\mathcal{Z}^\beta$ is a bounded operator if and only if (18) holds and

$$\sup_{n\geq 0}(n+1)^{\alpha-2}\|((g''\circ\varphi)(\varphi')^2+(g'\circ\varphi)\varphi'')\varphi^n\|_{v_\beta}\asymp\sup_{z\in D}\frac{(1-|z|^2)^\beta}{(1-|\varphi(z)|^2)^{\alpha-2}}\times|g''(\varphi(z))(\varphi'(z))^2+g'(\varphi(z))\varphi''(z)|<\infty. \tag{2.20}$$

Proof Suppose that $C_\varphi U_g$ is bounded from $\mathcal{Z}^\alpha$ to $\mathcal{Z}^\beta$.

(i) **Case** $0 < \alpha < 1$**.** Using functions $f = 1 \in \mathcal{Z}^\alpha$ and $f = z \in \mathcal{Z}^\alpha$, we obtain

$$\sup_{z\in D}(1-|z|^2)^\beta |g''(\varphi(z))(\varphi'(z))^2 + g'(\varphi(z))\varphi''(z)| < \infty, \tag{2.21}$$

and

$$\sup_{z\in D}(1-|z|^2)^\beta |g'(\varphi(z))(\varphi'(z))^2| < \infty. \tag{2.22}$$

Then we obtain that $((g'\circ\varphi)(\varphi'') + (g''\circ\varphi)(\varphi')^2) \in H^\infty_{v_\beta}$ and $(g'\circ\varphi)(\varphi')^2 \in H^\infty_{v_\beta}$ are necessary for all case.

For the converse implication, suppose that $((g'\circ\varphi)(\varphi'') + (g''\circ\varphi)(\varphi')^2) \in H^\infty_{v_\beta}$ and $(g'\circ\varphi)(\varphi')^2 \in H^\infty_{v_\beta}$. For $f \in \mathcal{Z}^\alpha$, it follows from Lemma 2.2 that

$$\begin{aligned}
(1-|z|^2)^\beta |(C_\varphi U_g f)''(z)| &= (1-|z|^2)^\beta |(\varphi'(z))^2 g'(\varphi(z)) f'(\varphi(z)) + (g''(\varphi(z))(\varphi'(z))^2 \\
&+ g'(\varphi(z))\varphi''(z))f(\varphi(z))| \le (1-|z|^2)^\beta |(\varphi'(z))^2 g'(\varphi(z)) f'(\varphi(z))| \\
&+ (1-|z|^2)^\beta |(g''(\varphi(z))(\varphi'(z))^2 + g'(\varphi(z))\varphi''(z))f(\varphi(z))| \\
&\le (1-|z|^2)^\beta |(\varphi'(z))^2 g'(\varphi(z))| \|f\|_{\mathcal{Z}^\alpha} \\
&+ (1-|z|^2)^\beta |g''(\varphi(z))(\varphi'(z))^2 + g'(\varphi(z))\varphi''(z)| \|f\|_{\mathcal{Z}^\alpha} \le C\|f\|_{\mathcal{Z}^\alpha},
\end{aligned}$$

$$\begin{aligned}
|C_\varphi U_g(f)(0)| &= \Big|\int_0^{\varphi(0)} f(\zeta) g'(\zeta) d\zeta\Big| \\
&\le \max_{\zeta\le|\varphi(0)|} |f(\zeta)| \max_{\zeta\le|\varphi(0)|} |g'(\zeta)| \le \max_{\zeta\le|\varphi(0)|} \|f\|_{\mathcal{Z}^\alpha} \max_{\zeta\le|\varphi(0)|} |g'(\zeta)|,
\end{aligned}$$

and

$$|(C_\varphi U_g f)'(0)| = |f(\varphi(0))\varphi'(0)g'(\varphi(0))| \le \|f\|_{\mathcal{Z}^\alpha} |\varphi'(0)||g'(\varphi(0))|.$$

Then $C_\varphi U_g$ is bounded. This completes the proof of (i).

(ii) **Case** $\alpha = 1$**.** We consider the test function $k_{\varphi(z)}(z)$ defined in (12) for every $z \in D$ with $|\varphi(z)| > \frac{1}{2}$. It follows that

$$\begin{aligned}
\|C_\varphi U_g k_{\varphi(z)}\|_{\mathcal{Z}^\beta} &\ge \sup_{z\in D}(1-|z|^2)^\beta |(C_\varphi U_g k_{\varphi(z)})''(z)| \\
&\ge \sup_{|\varphi(z)|>\frac{1}{2}} \log\Big(\frac{1}{1-|\varphi(z)|^2}\Big)(1-|z|^2)^\beta |(\varphi'(z))^2 g'(\varphi(z))| \\
&- \sup_{|\varphi(z)|>\frac{1}{2}} (1-|z|^2)^\beta |\varphi''(z)g'(\varphi(z)) + g''(\varphi(z))(\varphi(z))^2| |k_{\varphi(z)}(\varphi(z))|.
\end{aligned}$$

Since $((g'\circ\varphi)(\varphi'') + (g''\circ\varphi)(\varphi')^2) \in H^\infty_{v_\beta}$, and $\sup_{|\varphi(z)|>\frac{1}{2}} \|C_\varphi U_g k_{\varphi(z)}\|_{\mathcal{Z}_1} \le C$, we get

$$\begin{aligned}
&\sup_{|\varphi(z)|>\frac{1}{2}} \log\frac{1}{1-|\varphi(z)|^2}(1-|z|^2)^\beta |(\varphi'(z))^2 g'(\varphi(z))| \\
&\le \sup_{|\varphi(z)|>\frac{1}{2}} (1-|z|^2)^\beta |\varphi''(z)g'(\varphi(z)) + g''(\varphi(z))(\varphi(z))^2| |k_{\varphi(z)}(\varphi(z))| + \|C_\varphi U_g k_{\varphi(z)}\| < \infty.
\end{aligned}$$

Then we have

$$\begin{aligned}
& \sup_{z\in D}\log(\frac{1}{1-|\varphi(z)|^2})(1-|z|^2)^\beta|(\varphi'(z))^2g'(\varphi(z))| \\
\leq & \sup_{|\varphi(z)|\leq\frac{1}{2}}\log(\frac{1}{1-|\varphi(z)|^2})(1-|z|^2)^\beta|(\varphi'(z))^2g'(\varphi(z))| \\
+ & \sup_{|\varphi(z)|>\frac{1}{2}}\log(\frac{1}{1-|\varphi(z)|^2})(1-|z|^2)^\beta|(\varphi'(z))^2g'(\varphi(z))| \\
\leq & C+\log(\frac{4}{3})\|(g'\circ\varphi)(\varphi')^2\|_{v_\beta}<\infty.
\end{aligned}$$

On the other hand, from (3) and Lemma 2.4, we have

$$\sup_{n\geq 0}(\log n)\|(g'\circ\varphi)(\varphi')^2\varphi^n\|_{v_\beta}\asymp\sup_{z\in D}(1-|z|^2)^\beta\log(\frac{2}{1-|\varphi(z)|^2})|g'(\varphi(z))(\varphi'(z))^2|.$$

Hence (17) holds.

The converse implication can be shown as in the proof of (i).

(iii) **Case** $1<\alpha<2$**.** $((g'\circ\varphi)(\varphi'')+(g''\circ\varphi)(\varphi')^2)\in H^\infty_{v_\beta}$ has been proved above. We take the test function $f_{\varphi(z)}$ in (9) for every $z\in D$ with $|\varphi(z)|>\frac{1}{2}$, by the same way as (ii), we can obtain that (18) holds.

The converse implication can be shown as in the proof of (i).

(iv) **Case** $\alpha=2$**.** We have prove that (19) holds above. To prove (20), we consider another test function $t_a(z)=\log\frac{2}{1-\overline{a}z}$. Clearly $t_a\in\mathcal{Z}^2$ and $\sup_{\frac{1}{2}<|a|<1}\|t_a\|_{\mathcal{Z}^2}<\infty$. For every $z\in D$ with $|\varphi(z)|>\frac{1}{2}$, it follows that

$$\begin{aligned}
\sup_{z\in D}(1-|z|^2)^\beta|(C_\varphi U_g t_a)''(z)| \geq & \sup_{|\varphi(z)|>\frac{1}{2}}\log(\frac{1}{1-|\varphi(z)|^2})(1-|z|^2)^\beta|(\varphi'(z))^2g''(\varphi(z)) \\
& + g'(\varphi(z))\varphi''(z)|-\sup_{|\varphi(z)|>\frac{1}{2}}\frac{(1-|z|^2)^\beta}{1-|\varphi(z)|^2}|g'(\varphi(z))(\varphi(z))^2|.
\end{aligned}$$

Applying (19) we get

$$\begin{aligned}
& \sup_{|\varphi(z)|>\frac{1}{2}}\log(\frac{1}{1-|\varphi(z)|^2})(1-|z|^2)^\beta|(\varphi'(z))^2g''(\varphi(z))+g'(\varphi(z))\varphi''(z)| \\
\leq & \sup_{|\varphi(z)|>\frac{1}{2}}\frac{(1-|z|^2)^\beta}{1-|\varphi(z)|^2}|g'(\varphi(z))(\varphi(z))^2|+\|C_\varphi U_g t_a\|<\infty.
\end{aligned}$$

Noting $((g'\circ\varphi)(\varphi'')+(g''\circ\varphi)(\varphi')^2)\in H^\infty_{v_\beta}$, and using Lemma 2.4 and (3), we conclude that (20) holds.

(v) **Case** $\alpha>2$**.** We have proved that (18) holds above. Applying test function $f_{\varphi(z)}$ in (9) for every $z\in D$ with $|\varphi(z)|>\frac{1}{2}$, we have

$$\begin{aligned}
S_1=\sup_{|\varphi(z)|>\frac{1}{2}}(1-|z|^2)^\beta|\frac{(\varphi'(z))^2g'(\varphi(z))}{\overline{\varphi(z)}(1-|\varphi(z)|^2)^{\alpha-1}}| = & \sup_{|\varphi(z)|>\frac{1}{2}}(1-|z|^2)^\beta|(\varphi'(z))^2g'(\varphi(z))||f'_{\varphi(z)}(\varphi(z))| \\
\leq & \sup_{|\varphi(z)|>\frac{1}{2}}\|C_\varphi U_g(f_{\varphi(z)})\|_{\mathcal{Z}^\beta}<\infty.
\end{aligned}$$

With the same calculation for test function $t_{\varphi(z)}(\varphi(z)) = \frac{(1-|\varphi(z)|^2)^2}{(1-\overline{\varphi(z)}z)^\alpha}$ with $|\varphi(z)| > \frac{1}{2}$, then $\sup_{|\varphi(z)|>\frac{1}{2}} \|t_{\varphi(z)}\|_{\mathcal{Z}^\alpha} \leq C$, and we have that

$$\begin{aligned} S_2 &= \sup_{|\varphi(z)|>\frac{1}{2}} (1-|z|^2)^\beta \left|\frac{(\varphi'(z))^2 g''(\varphi(z)) + g'(\varphi(z))\varphi''(z)}{(1-|\varphi(z)|^2)^{\alpha-2}} + \alpha\overline{\varphi(z)}\frac{(\varphi'(z))^2 g'(\varphi(z))}{(1-|\varphi(z)|^2)^{\alpha-1}}\right| \\ &= \sup_{|\varphi(z)|>\frac{1}{2}} (1-|z|^2)^\beta |(C_\varphi U_g(t_{\varphi(z)}))''(\varphi(z))| \leq \|C_\varphi U_g\|_{\mathcal{Z}^\alpha \to \mathcal{Z}^\beta} \sup_{|\varphi(z)|>\frac{1}{2}} \|t_{\varphi(z)}\|_{\mathcal{Z}^\alpha} < \infty. \end{aligned}$$

Therefore,

$$\begin{aligned} &\sup_{|\varphi(z)|>\frac{1}{2}} (1-|z|^2)^\beta \left|\frac{(\varphi'(z))^2 g''(\varphi(z)) + g'(\varphi(z))\varphi''(z)}{(1-|\varphi(z)|^2)^{\alpha-2}}\right| \\ &\leq S_2 + \alpha \sup_{|\varphi(z)|>\frac{1}{2}} \frac{|(\varphi'(z))^2 g'(\varphi(z))|}{(1-|\varphi(z)|^2)^{\alpha-1}} \\ &\leq S_2 + \alpha \sup_{|\varphi(z)|>\frac{1}{2}} \frac{|(\varphi'(z))^2 g'(\varphi(z))|}{|\overline{\varphi(z)}|(1-|\varphi(z)|^2)^{\alpha-1}} \leq S_2 + \alpha S_1 < \infty. \end{aligned}$$

Since $((g'\circ\varphi)(\varphi'') + (g''\circ\varphi)(\varphi')^2) \in H^\infty_{v_\beta}$, we conclude that (21) holds.

Theorem 2.4 *Let φ be an analytic self-map of D and $g \in H(D)$.*

(i) If $0 < \alpha < 1$, then $U_g C_\varphi : \mathcal{Z}^\alpha \to \mathcal{Z}^\beta$ is a bounded operator if and only if $g'\varphi' \in H^\infty_{v_\beta}$ and $g'' \in H^\infty_{v_\beta}$.

(ii) If $\alpha = 1$, then $U_g C_\varphi : \mathcal{Z}^\alpha \to \mathcal{Z}^\beta$ is a bounded operator if and only if $g'' \in H^\infty_{v_\beta}$ and

$$\sup_{n\geq 0}(\log n)\|(g'\varphi')\varphi^n\|_{v_\beta} \asymp \sup_{z\in D}(1-|z|^2)^\beta \log\frac{2}{1-|\varphi(z)|^2}|g'(z)\varphi'(z)| < \infty. \tag{2.23}$$

(iii) If $1 < \alpha < 2$, then $U_g C_\varphi : \mathcal{Z}^\alpha \to \mathcal{Z}^\beta$ is a bounded operator if and only if $g'' \in H^\infty_{v_\beta}$ and

$$\sup_{n\geq 0}(n+1)^{\alpha-1}\|(g'\varphi')\varphi^n\|_{v_\beta} \asymp \sup_{z\in D}\frac{(1-|z|^2)^\beta}{(1-|\varphi(z)|^2)^{\alpha-1}}|g'(z)\varphi'(z)| < \infty. \tag{2.24}$$

(iv) If $\alpha = 2$, then $U_g C_\varphi : \mathcal{Z}^\alpha \to \mathcal{Z}^\beta$ is a bounded operator if and only if

$$\sup_{n\geq 0}(n+1)\|(g'\varphi')\varphi^n\|_{v_\beta} \asymp \sup_{z\in D}\frac{(1-|z|^2)^\beta}{(1-|\varphi(z)|^2)}|g'(z)\varphi'(z)| < \infty, \tag{2.25}$$

and

$$\sup_{n\geq 0}(\log n)\|(g'')\varphi^n\|_{v_\beta} \asymp \sup_{z\in D}(1-|z|^2)^\beta \log\frac{2}{(1-|\varphi(z)|^2)}|g''(z)| < \infty. \tag{2.26}$$

(v) If $\alpha > 2$, then $U_g C_\varphi : \mathcal{Z}^\alpha \to \mathcal{Z}^\beta$ is a bounded operator if and only if (25) holds and

$$\sup_{n\geq 0}(n+1)^{\alpha-2}\|(g'')\varphi^n\|_{v_\beta} \asymp \sup_{z\in D}\frac{(1-|z|^2)^\beta}{(1-|\varphi(z)|^2)^{\alpha-2}}|g''(z)| < \infty. \tag{2.27}$$

The proof is similar to that of Theorem 2.3, the details are omitted.

3 Essential norms

In this section we estimate the essential norms of these integral-type operators on *Zygmund* type spaces in terms of g, φ, their derivatives and the n-th power φ^n of φ.

Let $\widetilde{\mathcal{Z}^\alpha} = \{f \in \mathcal{Z}^\alpha : f(0) = f'(0) = 0\}$ and $\widetilde{\mathcal{B}^\alpha} = \{f \in \mathcal{B}^\alpha : f(0) = 0\}$. We note that every compact operator $T \in \mathbf{K}(\widetilde{\mathcal{Z}^\alpha}, \mathcal{Z}^\beta)$ can be extended to a compact operator $K \in \mathbf{K}(\mathcal{Z}^\alpha, \mathcal{Z}^\beta)$. In fact, for every $f \in \mathcal{Z}^\alpha$, $f - f(0) - f'(0)z \in \widetilde{\mathcal{Z}^\alpha}$, and we can define $K(f) = T(f - f(0) - f'(0)z) + f(0) + f'(0))z$.

For $r \in (0,1)$, we consider the compact operator $K_r : \mathcal{Z}^\alpha \to \mathcal{Z}^\beta$ defined by $K_r f(z) = f(rz)$.

Lemma 3.1 *If $X(V_gC_\varphi, C_\varphi V_g, C_\varphi U_g, U_gC_\varphi)$ is a bounded operator from $\mathcal{Z}^\alpha$ to $\mathcal{Z}^\beta$ space, then*

$$\|X\|_{e,\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} = \|X\|_{e,\widetilde{\mathcal{Z}^\alpha}\to\mathcal{Z}^\beta}.$$

Proof Clearly $\|X\|_{e,\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} \geq \|X\|_{e,\widetilde{\mathcal{Z}^\alpha}\to\mathcal{Z}^\beta}$. Then we prove $\|X\|_{e,\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} \leq \|X\|_{e,\widetilde{\mathcal{Z}^\alpha}\to\mathcal{Z}^\beta}$.

Let $T \in K(\mathcal{Z}^\alpha, \mathcal{Z}^\beta)$ be given. Let $\{r_n\}$ be an increasing sequence in $(0,1)$ converging to 1 and $\mathbf{A} = \{h | h = a + bz\}$, the closed subspace of $\mathcal{Z}^\alpha$. Then

$$\begin{aligned}
\|X - T\|_{\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} &= \sup_{f\in\mathcal{Z}^\alpha, \|f\|_{\mathcal{Z}^\alpha}\leq 1} \|X(f) - T(f)\|_{\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} \\
&\leq \sup_{f\in\mathcal{Z}^\alpha, \|f\|_{\mathcal{Z}^\alpha}\leq 1} \|X(f - f(0) - f'(0)z) - T(f - f(0) - f'(0)z)\|_{\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} \\
&+ \sup_{f\in\mathcal{Z}^\alpha, \|f\|_{\mathcal{Z}^\alpha}\leq 1} \|X(f(0) + f'(0)z) - T(f(0) + f'(0)z)\|_{\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} \\
&\leq \sup_{g\in\widetilde{\mathcal{Z}^\alpha}, \|g\|_{\mathcal{Z}^\alpha}\leq 1} \|X(g) - T|_{\widetilde{\mathcal{Z}^\alpha}}(g)\|_{\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} + \sup_{h\in\alpha, \|h\|_\alpha\leq 1} \|X(h) - T|_\alpha(h)\|_{\mathcal{Z}^\alpha\to\mathcal{Z}^\beta}.
\end{aligned}$$

Hence

$$\begin{aligned}
\inf_{T\in K(\mathcal{Z}^\alpha,\mathcal{Z}^\beta)} \|X - T\|_{\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} &\leq \inf_{T\in K(\mathcal{Z}^\alpha,\mathcal{Z}^\beta)} \|X - T|_{\widetilde{\mathcal{Z}^\alpha}}\|_{\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} + \inf_{T\in K(\mathcal{Z}^\alpha,\mathcal{Z}^\beta)} \|X - T|_\alpha\|_{\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} \\
&\leq \|X\|_{e,\widetilde{\mathcal{Z}^\alpha}\to\mathcal{Z}^\beta} + \lim_{n\to\infty} \|X(I - K_{r_n})\|_{\alpha\to\mathcal{Z}^\beta}.
\end{aligned}$$

Since $\lim_{n\to\infty} \|X(I - K_{r_n})\|_{\alpha\to\mathcal{Z}^\beta} = 0$, we have, $\|X\|_{e,\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} \leq \|X\|_{e,\widetilde{\mathcal{Z}^\alpha}\to\mathcal{Z}^\beta}$, and the proof is finished .

Let $D_\alpha : \mathcal{Z}^\alpha \to \mathcal{B}_\alpha$ and $S_\alpha : \mathcal{B}^\alpha \to H^\infty_{v_\alpha}$ be the derivative operators. Then clearly D_α and S_α are linear isometries on $\widetilde{\mathcal{Z}^\alpha}$ and $\widetilde{\mathcal{B}^\alpha}$, respectively and

$$S_\beta D_\beta V_g C_\varphi D_\alpha^{-1} S_\alpha^{-1} = g(\varphi')C_\varphi + g' C_\varphi S_\alpha^{-1},$$

Therefore

$$\|V_g C_\varphi\|_{e,\widetilde{\mathcal{Z}^\alpha}\to\mathcal{Z}^\beta} \leq \|g' C_\varphi\|_{e,\widetilde{\mathcal{B}^\alpha}\to H^\infty_{v_\beta}} + \|g(\varphi')C_\varphi\|_{e,H^\infty_{v_\alpha}\to H^\infty_{v_\beta}}. \tag{3.1}$$

Similarly, we have

$S_\beta D_\beta C_\varphi V_g D_\alpha^{-1} S_\alpha^{-1} = (g\circ\varphi(\varphi')^2)C_\varphi + (g'\circ\varphi(\varphi')^2 + g\circ\varphi(\varphi''))C_\varphi S_\alpha^{-1}$,

$$\|C_\varphi V_g\|_{e,\widetilde{\mathcal{Z}^\alpha}\to\mathcal{Z}^\beta} \leq \|((g'\circ\varphi(\varphi')^2))C_\varphi\|_{e,H^\infty_{v_\alpha}\to H^\infty_{v_\beta}} + \|((g'\circ\varphi)(\varphi')^2 + g\circ\varphi(\varphi''))C_\varphi\|_{e,\widetilde{\mathcal{B}^\alpha}\to H^\infty_{v_\beta}}. \tag{3.2}$$

$$S_\beta D_\beta C_\varphi U_g D_\alpha^{-1} S_\alpha^{-1} = (g'' \circ \varphi(\varphi')^2 + g' \circ \varphi(\varphi'')) C_\varphi D_\alpha^{-1} S_\alpha^{-1} + (g' \circ \varphi(\varphi')^2) C_\varphi S_\alpha^{-1},$$

$$\|C_\varphi U_g\|_{e,\widetilde{\mathcal{Z}^\alpha} \to \mathcal{Z}^\beta} \leq \|((g' \circ \varphi)(\varphi')^2) C_\varphi\|_{e,\widetilde{\mathcal{B}^\alpha} \to H^\infty_{v_\beta}} + \|((g'' \circ \varphi)(\varphi')^2 + g' \circ \varphi(\varphi'')) C_\varphi\|_{e,\widetilde{\mathcal{Z}^\alpha} \to H^\infty_{v_\beta}}. \quad (3.3)$$

$$S_\beta D_\beta U_g C_\varphi D_\alpha^{-1} S_\alpha^{-1} = g'' C_\varphi D_\alpha^{-1} S_\alpha^{-1} + g'(\varphi') C_\varphi S_\alpha^{-1},$$

$$\|U_g C_\varphi\|_{e,\widetilde{\mathcal{Z}^\alpha} \to \mathcal{Z}^\beta} \leq \|g'(\varphi')\|_{e,\widetilde{\mathcal{B}^\alpha} \to H^\infty_{v_\beta}} + \|g'' C_\varphi\|_{e,\widetilde{\mathcal{Z}^\alpha} \to H^\infty_{v_\beta}}. \quad (3.4)$$

Theorem 3.1 *Let φ be an analytic self-map of D and $g \in H(D)$, and $V_g C_\varphi : \mathcal{Z}^\alpha \to \mathcal{Z}^\beta$ is a bounded operator.*

(i) If $0 < \alpha < 1$, then

$$\|V_g C_\varphi\|_{e,\mathcal{Z}^\alpha \to \mathcal{Z}^\beta} \asymp \limsup_{n\to\infty} (n+1)^\alpha \|(g\varphi')\varphi^n\|_{v_\beta}. \quad (3.5)$$

(ii) If $\alpha = 1$, then

$$\|V_g C_\varphi\|_{e,\mathcal{Z} \to \mathcal{Z}^\beta} \asymp \max\{\limsup_{n\to\infty} (n+1)\|(g\varphi')\varphi^n\|_{v_\beta}, \limsup_{n\to\infty} (\log n)\|(g')\varphi^n\|_{v_\beta}\}. \quad (3.6)$$

(iii) If $\alpha > 1$, then

$$\|V_g C_\varphi\|_{e,\mathcal{Z}^\alpha \to \mathcal{Z}^\beta} \asymp \max\{\limsup_{n\to\infty} (n+1)^\alpha \|(g\varphi')\varphi^n\|_{v_\beta}, \limsup_{n\to\infty} (n+1)^{\alpha-1} \|(g')\varphi^n\|_{v_\beta}\}. \quad (3.7)$$

Proof (i) We start with the upper bound. First we show that $g'C_\varphi$ is a compact weighted composition operator for $\mathcal{B}^\alpha$ into $H^\infty_{v_\beta}$. Suppose that $\{f_n\}$ is bounded sequence in $\mathcal{B}^\alpha$. From Lemma 3.6 in [9], $\{f_n\}$ has a subsequence $\{f_{n_k}\}$ which converges uniformly on D to a function, which we can assume to be identically zero. Then it follows from Theorem 2.1 and Lemma 2.1 that

$$\lim_{k\to\infty} \sup_{z\in D} (1-|z|^2)^\beta |g'(z)||f_{n_k}(\varphi(z))| \leq C \lim_{k\to\infty} \sup_{z\in D} |f_{n_k}(z)| = 0,$$

which shows that $g'C_\varphi : \mathcal{B}^\alpha \to H^\infty_{v_\beta}$ is a compact operator and $\|g'C_\varphi\|_{e,\mathcal{B}^\alpha \to H^\infty_{v_\beta}} = 0$. Applying (29), Lemma 2.5 and 3.1, we get that

$$\begin{aligned}
\|V_g C_\varphi\|_{e,\mathcal{Z}^\alpha \to \mathcal{Z}^\beta} \leq \|g(\varphi')C_\varphi\|_{e,H^\infty_{v_\alpha} \to H^\infty_{v_\beta}} &= \limsup_{|\varphi(z)|\to 1} \frac{(1-|z|^2)^\beta}{(1-|\varphi(z)|^2)^\alpha} |g(z)\varphi'(z)| \\
&= \limsup_{n\to\infty} \frac{\|(g\varphi')\varphi^n\|_{v_\beta}}{\|z^n\|_{v_\alpha}} \\
&= (\frac{e}{2\alpha})^\alpha \limsup_{n\to\infty} (n+1)^\alpha \|(g\varphi')\varphi^n\|_{v_\beta}.
\end{aligned}$$

For the lower bound, let $\{z_n\} \subseteq D$ with $|\varphi(z_n)| > \frac{1}{2}$ and $|\varphi(z_n)| \to 1$ as $n \to \infty$. Taking $g_n = g_{\varphi(z_n)}$ defined in (11), we obtain that $\{g_n\}$ is bounded sequence in $\mathcal{Z}^\alpha_0$ converging to 0 uniformly on compact subset of D and $\sup_{n\in N} \|g_n\|_{\mathcal{Z}^\alpha} \leq C$. For every compact operator $T : \mathcal{Z}^\alpha \to \mathcal{Z}^\beta$,

$$\begin{aligned}
C\|V_g C_\varphi - T\|_{\mathcal{Z}^\alpha \to \mathcal{Z}^\beta} &\geq \limsup_{n\to\infty} \|V_g C_\varphi g_n\|_{\mathcal{Z}^\beta} - \limsup_{n\to\infty} \|T(g_n)\|_{\mathcal{Z}^\beta} \\
&\geq \alpha \limsup_{n\to\infty} (1-|z_n|^2)^\beta \frac{|g(z_n)\varphi'(z_n)|}{(1-|\varphi(z_n)|^2)^\alpha}.
\end{aligned}$$

Now we use (2) and Lemma 2.4 to obtain that

$$\begin{aligned}\|V_gC_\varphi\|_{e,\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} &\geq \|V_gC_\varphi - T\|_{\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} \geq \frac{\alpha}{C}\limsup_{|\varphi(z_n)|\to 1}|g(z_n)\varphi'(z_n)|\frac{(1-|z_n|^2)^\beta}{(1-|\varphi(z_n)|^2)^\alpha}\\ &= \frac{\alpha}{C}\limsup_{n\to\infty}\frac{\|(g\varphi')\varphi^n\|_{v_\beta}}{\|z^n\|_{v_\alpha}}\\ &= C(\frac{e}{2\alpha})^\alpha\limsup_{n\to\infty}(n+1)^\alpha\|(g\varphi')\varphi^n\|_{v_\beta}.\end{aligned}$$

Hence (33) holds.

(ii) The boundedness of V_gC_φ guarantees that $(g\varphi')C_\varphi : H^\infty_{v_1}\to H^\infty_{v_\beta}$ and $(g')C_\varphi : \mathcal{B}\to H^\infty_{v_\beta}$ are bounded weighted composition operators. Theorem 3.4 in [15] ensures that

$$\|g'C_\varphi\|_{e,\mathcal{B}\to H^\infty_{v_\beta}} \asymp \lim_{|\varphi(z)|\to 1}(1-|z|^2)^\beta|g'(z)|\log\frac{2}{1-|\varphi(z)|^2}.$$

Now we use Lemma 2.4, 2.5 and (29) to conclude that

$$\begin{aligned}\|V_gC_\varphi\|_{e,\mathcal{Z}\to\mathcal{Z}^\beta} &\leq \|g'C_\varphi\|_{e,\widetilde{\mathcal{B}}\to H^\infty_{v_\beta}} + \|g(\varphi')C_\varphi\|_{e,H^\infty_{v_1}\to H^\infty_{v_\beta}}\\ &\leq C\limsup_{n\to\infty}\frac{\|(g\varphi')\varphi^n\|_{v_\beta}}{\|z^n\|_{v_1}} + C\limsup_{n\to\infty}\frac{\|g'\varphi^n\|_{v_\beta}}{\|z^n\|_{v_{\log}}}\\ &= C\limsup_{n\to\infty}(\log n)\|g'\varphi^n\|_{v_\beta} + \frac{Ce}{2}\limsup_{n\to\infty}(n+1)\|(g\varphi')\varphi^n\|_{v_\beta}\\ &\leq C\max\{\limsup_{n\to\infty}(n+1)\|(g\varphi')\varphi^n\|_{v_\beta}, \limsup_{n\to\infty}(\log n)\|(g')\varphi^n\|_{v_\beta}\}.\end{aligned}$$

On the other hand, let $\{z_n\}$ be a sequence in D such that $|\varphi(z_n)| > \frac{1}{2}$ and $|\varphi(z_n)|\to 1$ as $n\to\infty$. Given

$$h_n(z) = \frac{h(\overline{\varphi(z_n)}z)}{\overline{\varphi(z_n)}}(\log\frac{2}{1-|\varphi(z_n)|^2})^{-1} - (\int_0^z \log^3\frac{2}{1-\overline{\varphi(z_n)}\omega}d\omega)(\log\frac{2}{1-|\varphi(z_n)|^2})^{-2}, \quad (3.8)$$

where $h(z) = (z-1)((1+\log\frac{2}{1-z})^2+1)$. From [6] we know that $\{h_n\}$ is a bounded sequence in $\mathcal{Z}^0_1$ which converges to zero uniformly on compact subsets of D, and

$$h_n''(\varphi(z_n)) = \frac{-\overline{\varphi(z_n)}}{1-|\varphi(z_n)|^2}, \qquad h_n'(\varphi(z_n)) = 0, \qquad \sup_n\|h_n\|_{\mathcal{Z}} < +\infty.$$

For every compact operator $T : \mathcal{Z}\to\mathcal{Z}^\beta$, we have $\|T(h_n)\|_{\mathcal{Z}^\beta}\to 0$ as $n\to\infty$. By Lemma 2.4 and 2.5, we obtain that

$$\begin{aligned}C\|V_gC_\varphi - T\|_{\mathcal{Z}\to\mathcal{Z}^\beta} &\geq \|V_gC_\varphi h_n\|_{\mathcal{Z}\to\mathcal{Z}^\beta} \geq \limsup_{n\to\infty}(1-|z_n|^2)^\beta||g(z_n)\varphi'(z_n)|\frac{|\varphi(z_n)|}{1-|\varphi(z_n)|^2}\\ &\geq \limsup_{n\to\infty}\frac{\|(g\varphi')\varphi^n\|_{v_\beta}}{\|z^n\|_{v_1}}\\ &= \frac{e}{2}\limsup_{n\to\infty}(n+1)\|(g\varphi')\varphi^n\|_{v_\beta}.\end{aligned}$$

Now we take another function

$$f_n = \frac{h(\overline{\varphi(z_n)}z)}{\overline{\varphi(z_n)}}(\log\frac{2}{1-|\varphi(z_n)|^2})^{-1}. \tag{3.9}$$

From [6] we know that $\{f_n\}$ is a bounded sequence in $\mathcal{Z}_1^0$ which converges to zero uniformly on compact subsets of D, and $\sup_{n\geq 1}\|f_n\|_{\mathcal{Z}} < +\infty$. It follows from Lemma 2.4 and 2.5 that

$$\begin{aligned} C\|V_gC_\varphi\|_{e,\mathcal{Z}\to\mathcal{Z}^\beta} &\geq \lim_{n\to\infty}\sup\|V_gC_\varphi f_n\|_{\mathcal{Z}^\beta} \geq \limsup_{n\to\infty}(1-|z_n|^2)^\beta|g'(z_n)|\log\frac{2}{1-|\varphi(z_n)|^2} \\ &- \limsup_{n\to\infty}(1-|z_n|^2)^\beta|(g(z_n)\varphi'(z_n)|\frac{|\varphi(z_n)|}{1-|\varphi(z_n)|^2}. \end{aligned}$$

Noting that $\limsup_{n\to\infty}(n+1)\|(g\varphi')\varphi^n\|_{v_\beta} \leq \frac{2C}{e}\|V_gC_\varphi\|_{e,\mathcal{Z}\to\mathcal{Z}^\beta}$, we obtain

$$\begin{aligned} (C+\frac{2C}{e})\|V_gC_\varphi\|_{e,\mathcal{Z}\to\mathcal{Z}^\beta} &\geq \limsup_{|\varphi(z_n)|\to 1}(1-|z_n|^2)^\beta|g'(z_n)|\log\frac{2}{1-|\varphi(z_n)|^2} \\ &= \limsup_{n\to\infty}(\log n)\|(g')\varphi^n\|_{v_\beta}. \end{aligned}$$

Hence we have $\|V_gC_\varphi\|_{e,\mathcal{Z}\to\mathcal{Z}^\beta} \geq C\max\{\limsup_{n\to\infty}(n+1)\|(g\varphi')\varphi^n\|_{v_\beta}, \limsup_{n\to\infty}(\log n)\|(g')\varphi^n\|_{v_\beta}\}$.

(iii) Let $\alpha > 1$. The proof of the upper bound is similar to that of (ii). From the proof of (i), we get that for some constant C ,

$$C\|V_gC_\varphi\|_{e,\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} \geq \limsup_{|\varphi(z)|\to 1}|g(z)\varphi'(z)|\frac{(1-|z_n|^2)^\beta}{(1-|\varphi(z)|^2)^\alpha}. \tag{3.10}$$

Now, let $\{z_n\}$ be as before and note that the function $f_n = f_{\varphi(z_n)}$ given in (9). Then $\{f_n\}$ is bounded sequence in $\mathcal{Z}_0^\alpha$ converging to zero uniformly on compact subsets of D , therefore

$$\begin{aligned} C\|V_gC_\varphi\|_{e,\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} &\geq \lim_{n\to\infty}\|V_gC_\varphi(f_n)\|_{\mathcal{Z}^\beta} \\ &\geq 2\alpha\limsup_{n\to\infty}\frac{(1-|z_n|^2)^\beta}{(1-|\varphi(z_n)|^2)^{\alpha-1}}|g'(z_n)| - \limsup_{n\to\infty}\frac{(1-|z_n|^2)^\beta|g(z_n)\varphi'(z_n)|}{|\varphi(z_n)|(1-|\varphi(z_n)|^2)^\alpha}. \end{aligned}$$

By (38), we have

$$C\|V_gC_\varphi\|_{e,\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} \geq \limsup_{n\to\infty}\frac{(1-|z_n|^2)^\beta}{(1-|\varphi(z_n)|^2)^{\alpha-1}}|g'(z_n)|,$$

and the rest of the proof is similar to that of the previous, we omit it .

Theorem 3.2 *Let φ be an analytic self-map of D and $g \in H(D)$, and $C_\varphi U_g : \mathcal{Z}^\alpha \to \mathcal{Z}^\beta$ is a bounded operator.*

(i) If $0 < \alpha < 1$, then

$$\|C_\varphi U_g\|_{e,\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} = 0. \tag{3.11}$$

(ii) If $\alpha = 1$, then

$$\|C_\varphi U_g\|_{e,\mathcal{Z}\to\mathcal{Z}^\beta} \asymp \lim_{n\to\infty} \sup(\log n)\|((g'\circ\varphi)(\varphi')^2)\varphi^n\|_{v_\beta}. \tag{3.12}$$

(iii) If $1 < \alpha < 2$, then

$$\|C_\varphi U_g\|_{e,\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} \asymp \lim_{n\to\infty} \sup(n+1)^{\alpha-1}\|((g'\circ\varphi)(\varphi')^2)\varphi^n\|_{v_\beta}. \tag{3.13}$$

(iv) If $\alpha = 2$, then

$$\begin{aligned}\|C_\varphi U_g\|_{e,\mathcal{Z}^2\to\mathcal{Z}^\beta} \asymp \max\{&\lim_{n\to\infty} \sup(n+1)\|((g'\circ\varphi)(\varphi')^2)\varphi^n\|_{v_\beta},\\ &\lim_{n\to\infty} \sup(\log n)\|((g''\circ\varphi)(\varphi')^2 + (g'\circ\varphi)(\varphi''))\varphi^n\|_{v_\beta}\}.\end{aligned} \tag{3.14}$$

(v) If $\alpha > 2$, then

$$\begin{aligned}\|C_\varphi U_g\|_{e,\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} \asymp \max\{&\lim_{n\to\infty} \sup(n+1)^{\alpha-1}\|((g'\circ\varphi)(\varphi')^2)\varphi^n\|_{v_\beta},\\ &\lim_{n\to\infty} \sup(n+1)^{\alpha-2}\|((g''\circ\varphi)(\varphi')^2 + (g'\circ\varphi)(\varphi''))\varphi^n\|_{v_\beta}\}.\end{aligned} \tag{3.15}$$

Proof (i) For the compactness of $((g'\circ\varphi)(\varphi')^2)C_\varphi$, the arguement is similar to the proof of Theorem 3.1(i), then we have $\|((g'\circ\varphi)(\varphi')^2)C_\varphi\|_{e,\mathcal{B}^\alpha\to H^\infty_{v_\beta}} = 0$. Hence by (31) and Lemma 2.3, we get that $\|C_\varphi U_g\|_{e,\mathcal{Z}^\alpha\to\mathcal{Z}^\beta} = 0$.

Next we will prove (ii). The boundedness of $C_\varphi U_g$ guarantees that $((g'\circ\varphi)(\varphi')^2)C_\varphi : \mathcal{B}\to H^\infty_{v_\beta}$ and $((g''\circ\varphi)(\varphi')^2 + (g'\circ\varphi)(\varphi''))C_\varphi : \mathcal{Z}\to H^\infty_{v_\beta}$ are bounded weighted composition operators. We know that if $uC_\varphi : \mathcal{Z}\to H^\infty_{v_\beta}$ is a bounded operator, then uC_φ is a compact operator by Lemma 2.3. Hence we consider the boundedness of $C_\varphi U_g : \mathcal{Z}\to\mathcal{Z}^\beta$ just consider that $((g'\circ\varphi)(\varphi')^2)C_\varphi : \mathcal{B}\to H^\infty_{v_\beta}$ is a bounded operator.

Theorem 3.4 in ([15]) ensures that

$$\|((g'\circ\varphi)(\varphi')^2)C_\varphi\|_{e,\mathcal{B}\to H^\infty_{v_\beta}} \asymp \lim_{|\varphi|\to 1}(1-|z|^2)^\beta|g'(\varphi(z))(\varphi'(z))^2|\log\frac{2}{1-|\varphi(z)|^2},$$

From (31), Lemma 2.4 and 2.5, we have

$$\begin{aligned}\|C_\varphi U_g\|_{e,\mathcal{Z}\to\mathcal{Z}^\beta} &\le C\limsup_{n\to\infty}\frac{\|((g'\circ\varphi)(\varphi')^2)\varphi^n\|_{v_\beta}}{\|z^n\|_{v_{\log}}}\\ &= C\limsup_{n\to\infty}(\log n)\|((g'\circ\varphi)(\varphi')^2)\varphi^n\|_{v_\beta}\end{aligned}$$

In order to prove $\|C_\varphi U_g\|_{e,\mathcal{Z}\to\mathcal{Z}^\beta} \ge \limsup_{n\to\infty}(\log n)\|((g'\circ\varphi)(\varphi')^2\}$, we take the function

$$g_n(z) = \frac{\overline{\varphi(z_n)}z-1}{\overline{\varphi(z_n)}}\big((1+\log\frac{1}{1-\overline{\varphi(z_n)}z})^2+1\big)(\log\frac{1}{1-|\varphi(z_n)|^2})^{-1} - a_n, \tag{3.16}$$

where

$$a_n = \frac{|\varphi(z_n)|^2-1}{\overline{\varphi(z_n)}}\big((1+\log\frac{1}{1-|\varphi(z_n)|^2})^2+1\big)(\log\frac{1}{1-|\varphi(z_n)|^2})^{-1},$$

and $\lim_{n\to\infty} a_n = 0$. From [18] we obtain that $\{g_n\}$ is a bounded sequence in $\mathcal{Z}_1^0$ which converges to zero uniformly on compact subsets of D. By a directly calculation, we have

$$g_n(\varphi(z_n)) = 0, \quad g_n'(\varphi(z_n)) = \log\frac{1}{1-|\varphi(z_n)|^2}.$$

For every compact operator $T : \mathcal{Z} \to \mathcal{Z}^\beta$, we have $\|T(h_n)\|_{\mathcal{Z}^\beta} \to 0$ as $n \to \infty$. Let $M = \sup_{n\geq 1}\|g_n\|_{\mathcal{Z}^\beta}$. It follows from Lemma 2.5 that

$$\begin{aligned} M\|C_\varphi U_g - T\|_{e,\mathcal{Z}\to\mathcal{Z}^\beta} &\geq \|C_\varphi U_g f_n\|_{e,\mathcal{Z}\to\mathcal{Z}^\beta} \\ &\geq \limsup_{n\to\infty}(1-|z_n|^2)^\beta |g'(\varphi(z_n))(\varphi'(z_n))^2| \log\frac{1}{1-|\varphi(z_n)|^2} \\ &\geq \limsup_{n\to\infty}\frac{\|((g'\circ\varphi)(\varphi')^2)\varphi^n\|_{v_\beta}}{\|z^n\|_v} \\ &\geq \limsup_{n\to\infty}(\log n)\|((g'\circ\varphi)(\varphi')^2)\varphi^n\|_{v_\beta}. \end{aligned}$$

This completes the proof.

The proof of (iii) is the same as that of Theorem 5(iii), we don't prove it again.

(iv) Let $\alpha = 2$. Applying Lemma 2.3 (ii) and Theorem 3.2 in [16], we get that

$$\|(g'\circ\varphi)(\varphi')^2 C_\varphi\|_{e,\mathcal{B}^2\to H^\infty_{v_\beta}} \asymp \limsup_{n\to\infty}(n+1)\|(g'\circ\varphi)(\varphi')^2\varphi^n\|_{v_\beta} \tag{3.17}$$

and

$$\|((g''\circ\varphi)(\varphi')^2+(g'\circ\varphi)\varphi'')C_\varphi\|_{e,\mathcal{Z}_2\to H^\infty_{v_\beta}} \asymp \limsup_{n\to\infty}(\log n)\|((g''\circ\varphi)(\varphi')^2+(g'\circ\varphi)\varphi'')\varphi^n\|_{v_\beta}, \tag{3.18}$$

which yields the upper bound by (31).

With the same arguements as in the proof of Theorem 2.3 and 3.1, for some constant C, we have

$$C\|C_\varphi U_g\|_{e,\mathcal{Z}^2\to\mathcal{Z}^\beta} \geq \limsup_{n\to\infty}(n+1)\|((g'\circ\varphi)(\varphi')^2)\varphi^n\|_{v_\beta}.$$

On the other hand, let $\{z_n\} \subseteq D$ with $|\varphi(z_n)| > \frac{1}{2}$ and $|\varphi(z_n)| \to 1$ as $n \to \infty$. Let the test function

$$O_n(z) = (1+(\log\frac{2}{1-\overline{\varphi(z_n)}z})^2)(\log\frac{2}{1-|\varphi(z_n)|^2})^{-1}.$$

From [1] we obtain that $\{O_n\}$ is a bounded sequence in $\mathcal{Z}_0^2$ which converges to zero uniformly on compact subsets of D, and

$$\begin{aligned} \lim_{n\to\infty}(1-|z_n|^2)^\beta |g'(\varphi(z_n))(\varphi'(z_n))^2||O_n'(\varphi(z_n))| &= 2\lim_{n\to\infty}\frac{(1-|z_n|^2)^\beta}{1-|\varphi(z_n)|^2}|g'(\varphi(z_n))(\varphi'(z_n))^2||\varphi(z_n)| \\ &\leq C\|C_\varphi U_g\|_{e,\mathcal{Z}^2\to\mathcal{Z}^\beta}. \end{aligned}$$

Applying Theorem 2.3 we get

$$\begin{aligned} C\|C_\varphi U_g\|_{e,\mathcal{Z}_2\to\mathcal{Z}^\beta} &\geq \|C_\varphi U_g(O_n)\|_{\mathcal{Z}^\beta} \\ &\geq \lim_{n\to\infty}(1-|z_n|^2)^\beta |g''(\varphi(z_n))(\varphi'(z_n))^2 + g'(\varphi(z_n))\varphi''(z_n)||O_n(\varphi(z_n))| \\ &- \lim_{n\to\infty}(1-|z_n|^2)^\beta |g'(\varphi(z_n))(\varphi'(z_n))^2||O_n'(\varphi(z_n))| \\ &\geq \lim_{n\to\infty}(1-|z_n|^2)^\beta |g''(\varphi(z_n))(\varphi'(z_n))^2 + g'(\varphi(z_n))\varphi''(z_n)|\log\frac{2}{1-|\varphi(z_n)|^2} \\ &- 2C\|C_\varphi U_g\|_{e,\mathcal{Z}^2\to\mathcal{Z}^\beta}. \end{aligned}$$

Hence

$$\lim_{n\to\infty}(1-|z_n|^2)^\beta|g''(\varphi(z_n))(\varphi'(z_n))^2+g'(\varphi(z_n))\varphi''(z_n)|\log\frac{2}{1-|\varphi(z_n)|^2}\le C\|C_\varphi U_g\|_{e,\mathcal{Z}^2\to\mathcal{Z}^\beta}.$$

On the other hand, the lower bound can be easily proved by Lemma 2.4 and 2.5.

If $\alpha>2$, the proof is similar to that of (iv) except that we now choose the test function $t_n(z)=\dfrac{(1-|\varphi(z_n)|^2)^2}{(1-\overline{\varphi(z_n)}z)^\alpha}$ instead of $O_n(z)$. This completes the proof of Theorem 3.2.

Using the same methods of Theorem 3.1 and 3.2, we can have the following results.

Theorem 3.3 *Let φ be an analytic self-map of D and $g\in H(D)$, and $C_\varphi V_g:\mathcal{Z}^\alpha\to\mathcal{Z}^\beta$ is a bounded operator.*

(i) If $0<\alpha<1$, then

$$\|C_\varphi V_g\|_{e,\mathcal{Z}^\alpha\to\mathcal{Z}^\beta}\asymp\limsup_{n\to\infty}(n+1)^\alpha\|((g\circ\varphi)(\varphi')^2)\varphi^n\|_{v_\beta}. \tag{3.19}$$

(ii) If $\alpha=1$, then

$$\begin{aligned}\|C_\varphi V_g\|_{e,\mathcal{Z}\to\mathcal{Z}^\beta}\asymp\max\{&\limsup_{n\to\infty}(n+1)\|((g\circ\varphi)(\varphi')^2)\varphi^n\|_{v_\beta},\\ &\limsup_{n\to\infty}(\log n)\|((g'\circ\varphi)(\varphi')^2+(g\circ\varphi)(\varphi''))\varphi^n\|_{v_\beta}\}.\end{aligned} \tag{3.20}$$

(iii) If $\alpha>1$, then

$$\begin{aligned}\|C_\varphi V_g\|_{e,\mathcal{Z}^\alpha\to\mathcal{Z}^\beta}\asymp\max\{&\limsup_{n\to\infty}(n+1)^\alpha\|((g\circ\varphi)(\varphi')^2)\varphi^n\|_{v_\beta},\\ &\limsup_{n\to\infty}(n+1)^{\alpha-1}\|((g'\circ\varphi)(\varphi')^2+(g\circ\varphi)(\varphi''))\varphi^n\|_{v_\beta}\}.\end{aligned} \tag{3.21}$$

Theorem 3.4 *Let φ be an analytic self-map of D and $g\in H(D)$, and $U_gC_\varphi:\mathcal{Z}^\alpha\to\mathcal{Z}^\beta$ is a bounded operator.*

(i) If $0<\alpha<1$, then

$$\|U_gC_\varphi\|_{e,\mathcal{Z}^\alpha\to\mathcal{Z}^\beta}=0. \tag{3.22}$$

(ii) If $\alpha=1$, then

$$\|U_gC_\varphi\|_{e,\mathcal{Z}\to\mathcal{Z}^\beta}\asymp\limsup_{n\to\infty}(\log n)\|((g'\varphi')\varphi^n\|_{v_\beta}\}. \tag{3.23}$$

(iii) If $1<\alpha<2$, then

$$\|U_gC_\varphi\|_{e,\mathcal{Z}^\alpha\to\mathcal{Z}^\beta}\asymp\limsup_{n\to\infty}(n+1)^{\alpha-1}\|((g'\varphi')\varphi^n\|_{v_\beta}\}. \tag{3.24}$$

(iv) If $\alpha=2$, then

$$\|U_gC_\varphi\|_{e,\mathcal{Z}^2\to\mathcal{Z}^\beta}\asymp\max\{\limsup_{n\to\infty}(n+1)\|((g'\varphi'))\varphi^n\|_{v_\beta},\limsup_{n\to\infty}(\log n)\|(g'')\varphi^n\|_{v_\beta}\}. \tag{3.25}$$

(v) If $\alpha>2$, then

$$\|U_gC_\varphi\|_{e,\mathcal{Z}^\alpha\to\mathcal{Z}^\beta}\asymp\max\{\limsup_{n\to\infty}(n+1)^{\alpha-1}\|((g'\varphi'))\varphi^n\|_{v_\beta},\limsup_{n\to\infty}(n+1)^{\alpha-2}\|(g'')\varphi^n\|_{v_\beta}\}. \tag{3.26}$$

Conflict of Interests

The authors declare that they have no conflict of interest.

References

[1] Esmaeili K, Lindström M. Weighted composition operators between Zygmund type spaces and their essential norms. Intege Equ Oper Theory, 2013, 75: 473-490

[2] Hyväarinen O, Kemppainen M, Lindström M, Rautio A, Saukko E. The essential norms of weighted composition operators on weighted Banach spaces of analytic function. Integr Equ Oper Theory, 2012, 72: 151C157

[3] Hyväarinen O, Lindström M. Estimates of essential norms of weighted com-position operators between Bloch type spaces. J Math AnalAppl, 2012, 393: 38C44

[4] Li S, Stevi̇c S. Weighted composition operators from Zygmund spaces into Bloch spaces. Appl Math Comput. 2008, 206: 825C831

[5] Li S, Stevi̇c S. Products of Volterra type operator and composition operator from H^∞ and Bloch spaces to Zygmund spaces. J Math Anal Appl. 2008, 345: 40C52

[6] Li S, Stevi̇c S. Generalized composition operators on Zygmund spaces and Bloch type spaces. J Math Anal Appl, 2008, 338: 1282C1295

[7] Li S, Stevi̇c S. Products of integral-type operators and composition operators between Bloch-type spaces. J Math Anal Appl, 2009, 349: 596C610

[8] Montes-Rodríguez A. Weighted composition operators on weighted Banach spaces of analytic functions. J London Math Soc, 2000, 61: 872C884

[9] Ohno S, Stroethoff K, Zhao R. Weighted composition operators between Bloch-type spaces. Rocky Mountain J Math, 2003, 33: 191C215

[10] Pommerenke C. Schlichte funktionen und analytische funktionen von beschränkter mittlerer Oszillation. Commentarii Mathematici Helvetici, 1997, 52: 591C602

[11] Stevi̇c S, Sharma A. Composition operators from weighted Bergman-Privalov spaces to Zygmund type spaces on the unit disk. Ann Polon Math, 2012, 105: 77C86

[12] Stevi̇c S. On an integral-type operator from Zygmund-type spaces to mixed-norm spaces on the unit ball. Abstr Appl Anal, 2010, Art. ID 198608, 7 pp.

[13] Stevi̇c S. Composition operators from the Hardy space to the Zygmund-type space on the upper half-plane. Abstr Appl Anal, 2009, Art. ID 161528, 8 pp.

[14] Stevi̇c S. On an integral operator from the Zygmund space to the Bloch-type space on the unit ball. Glasg Math J, 2009, 51: 275C287

[15] Stevi̇c S. Essential norms of weighted composition operators from the α-Bloch space to a weighted-type space on the unit ball. Abstr Appl Anal, 2008, Art. ID 279691

[16] Stevi̇c S. Weighted composition operators from Bergman-Privalov-type spaces to weighted-type spaces on the unit ball. Appl Math Comput, 2010, 217: 1939-1943

[17] Wolf E. Products of Volterra type operators and composition operators between weighted Bergman spaces of infinite order and weighted Bloch type spaces. Georgian Math J, 2010, 17: 621C627

[18] Ye S, Hu Q. Weighted composition operators on the Zygmund space. Abstr Appl Anal, 2012, Art.ID 462482, 18 pages.

[19] Ye S, Zhuo Z. Weighted composition operators from Hardy to Zygmund type spaces. Abstr Appl Anal, 2013, Art. ID 365286, 10 pages.

[20] Ye S. Products of Volterra-type operators and composition operators on logarithmic Bloch space. WSEAS Trans Math, 2013, 12: 180-188

[21] Ye S, Lin C. Composition followed by differentiation on the Zygmund space. Acta Math. Sinica, Chin. Ser., 2016, 59(1): 11C20.